

OPERADS AND ALGEBRAIC HOMOTOPY II: SUSPENSIONS

JUSTIN R. SMITH

1. INTRODUCTION

The present paper forms a continuation of [5]. Since that paper has not appeared, we begin by summarizing some of its relevant results.

The easiest way to convey the flavor of this paper's results is with a simple example. Suppose X and Y are pointed, simply-connected, 2-reduced simplicial sets. There are many topological invariants associated with the chain-complexes of X and Y — including the coproduct and S_n -equivariant *higher coproducts* (used to define Steenrod operations):

$$\begin{aligned} RS_n \otimes C(X) &\rightarrow C(X)^n \\ RS_n \otimes C(Y) &\rightarrow C(Y)^n \end{aligned}$$

for all $n > 1$, where:

1. RS_n is the bar-resolution of $\mathbb{Z}$ over $\mathbb{Z}S_n$.
2. $(*)^n$ denotes the n -fold tensor product over $\mathbb{Z}$ (with S_n acting by permuting factors).

Now suppose we know (from a purely *algebraic* analysis of these *chain-complexes*) that there exists a chain-map inducing homology isomorphisms

$$f: C(X) \rightarrow C(Y)$$

and making

$$(1.1) \quad \begin{array}{ccc} RS_n \otimes C(X) & \longrightarrow & C(X)^n \\ 1 \otimes f \downarrow & & \downarrow f^n \\ RS_n \otimes C(Y) & \longrightarrow & C(Y)^n \end{array}$$

commute for all $n > 1$ (requiring exact commutativity is unnecessarily restrictive, but we assume it to simplify this discussion).

Then our theory asserts that X and Y are homotopy equivalent via a geometric map inducing a map chain-homotopic to f . In fact, our theory goes further than this: if f is *any* chain-map making 1.1 commute for all $n > 1$, there exists a map of topological realizations

$$F: |X| \rightarrow |Y|$$

inducing a map equivalent to f (see [5, theorems 9.5 and 9.7] for the exact statements). Essentially, Steenrod operations on the *chain-level* determine:

- the homotopy types of spaces and maps,

1991 *Mathematics Subject Classification.* 55R91; Secondary: 18G30.

Key words and phrases. stable homotopy type, minimal models, operads.

- all obstructions to topologically realizing chain-maps.

I view this work as a generalization of Quillen's characterization of rational homotopy theory in [4] — he showed that rational homotopy types are determined by a commutative coalgebra structures on chain-complexes; we show that integral homotopy types are determined by coalgebra structures augmented by higher diagonals.

In [5, § 2], we define a category of m -coalgebras, $\mathfrak{L}_0$ and a localization of it, $\mathfrak{L}$ by a class of maps called elementary equivalences. This gives rise to a *homotopy theory* of m -coalgebras in terms of which we can state this paper's main result:

Theorem [5]: The functor

$$\mathcal{C}(*): \underline{\text{Homotop}}_0 \rightarrow \mathfrak{L}^+$$

defines an equivalence of categories and homotopy theories (in the sense of [3]), where $\underline{\text{Homotop}}_0$ is the homotopy category of pointed, simply-connected CW-complexes and continuous maps and $\mathfrak{L}^+ \subset \mathfrak{L}$ is the subcategory of topologically realizable m -coalgebras. In addition, there exists an equivalence of categories and homotopy theories

$$\mathcal{C}(*): \mathcal{F} \rightarrow \mathfrak{F}^+$$

where $\mathcal{F}$ is the homotopy category of finite, pointed, simply-connected simplicial sets and $\mathfrak{F}^+$ is the homotopy category of finitely generated, topologically realizable m -coalgebras in $\mathfrak{L}_0$ localized with respect to finitely generated equivalences in $\mathfrak{L}_0$.

In the spirit of our initial statement, one corollary to this result is

Corollary 9.20 of [5]: Let X and Y be pointed, simply-connected semisimplicial sets and let

$$f: \mathcal{C}(X) \rightarrow \mathcal{C}(Y)$$

be a chain-map between canonical chain-complexes. Then f is topologically realizable if and only if there exists an m -coalgebra C over $\mathfrak{S}$ and a factorization $f = f_\beta \circ f_\alpha$

$$\mathcal{C}(X_1) \xrightarrow{f_\alpha} C \xrightarrow[\iota]{f_\beta} \mathcal{C}(X_2)$$

where f_α is a morphism of m -coalgebras, ι is an elementary equivalence — an injection of m -coalgebras with acyclic, $\mathbb{Z}$ -free cokernel — and f_β is a chain map that is a left inverse to ι . If X and Y are finite, we may require C to be finitely generated.

The present paper will study *stable homotopy theory* using similar techniques.

In § 3, we define the important concept of suspension of operads and characterize suspensions of spaces in terms of m -coalgebras. This leads to the definition of the operad $\mathfrak{S}_\infty$, which can be thought of as an infinite desuspension of the operad $\mathfrak{S}$. The main result of this section is:

Corollary 3.24: Let X and Y be pointed, simply-connected semisimplicial sets. Then there exists an integer $k \geq 0$ and a pointed map of topological realizations $f: |S^k X| \rightarrow |S^k Y|$ if and only if there exists a morphism $g: \mathfrak{p}_0^* \mathcal{C}(X)^+ \rightarrow \mathfrak{p}_0^* \mathcal{C}(Y)^+$ in

$\Sigma^{-\infty}\mathfrak{L}$ (the category of coalgebras over $\mathfrak{S}_{-\infty}$). In this case, the following diagram commutes in $\mathfrak{L}$

$$(1.2) \quad \begin{array}{ccc} \Sigma^k(\mathfrak{U}^k)^*\mathcal{C}(X)^+ & \xrightarrow{\Sigma^k g} & \Sigma^k(\mathfrak{U}^k)^*\mathcal{C}(Y)^+ \\ \parallel & & \parallel \\ \mathcal{C}(S^k X)^+ & & \mathcal{C}(S^k Y)^+ \\ \downarrow \iota_{S^k X} & & \downarrow \iota_{S^k Y} \\ \mathcal{C}(\dot{\Delta}(|S^k X|)) & \xrightarrow{\mathcal{C}(\dot{\Delta}(f))} & \mathcal{C}(\dot{\Delta}(|S^k Y|)) \end{array}$$

where

1. Σ^k denotes suspension of chain-complexes and base-operads. This is suspension in the sense of dimension-shifting.
2. $\dot{\Delta}(\ast)$ denotes the 2-reduced singular complex functor.
3. $|\ast|$ denotes topological realization functor.
4. $\iota_X: \mathcal{C}(X) \rightarrow \mathcal{C}(\dot{\Delta}(|X|))$ is the canonical (injective) equivalence.

In particular X and Y are stably homotopy equivalent if and only if $\mathfrak{p}_0^*\mathcal{C}(X)^+$ and $\mathfrak{p}_0^*\mathcal{C}(Y)^+$ are equivalent in the category $\Sigma^{-\infty}\mathfrak{L}$.

In future papers, we will apply these results to Boardman's *stable homotopy category*. The stable homotopy category contains much more than suspensions of spaces — for instance, it also includes arbitrary “formal desuspensions” of spaces and “idealized” space-objects that characterize generalized homology theories. Even *spaces* behave differently in this category — as though already “infinitely suspended.” The objects of this category are *spectra* — sequences of spaces with maps from the suspension of each term of the sequence to the next term.

2. DEFINITIONS

In this section, we recall some of the relevant definitions from [5].

Definition 2.1. Let C and D be two graded $\mathbb{Z}$ -modules. A map of graded modules $f: C_i \rightarrow D_{i+k}$ will be said to be of degree k .

Remark 2.2. For instance the *differential* of a chain-complex will be regarded as a degree -1 map.

We will make extensive use of the Koszul Convention (see [1]) regarding signs in homological calculations:

Definition 2.3. If $f: C_1 \rightarrow D_1$, $g: C_2 \rightarrow D_2$ are maps, and $a \otimes b \in C_1 \otimes C_2$ (where a is a homogeneous element), then $(f \otimes g)(a \otimes b)$ is defined to be $(-1)^{\deg(g) \cdot \deg(a)} f(a) \otimes g(b)$.

Remark 2.4. This convention simplifies many of the common expressions that occur in homological algebra — in particular it eliminates complicated signs that occur in these expressions. For instance the differential, $\partial_{\otimes}$, of the tensor product $\partial_C \otimes 1 + 1 \otimes \partial_D$.

Throughout this entire paper we will follow the convention that group-elements act on the left. Multiplication of elements of symmetric groups will be carried out accordingly — i.e.

$$\begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & 1 & 4 \end{pmatrix} \cdot \begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 3 & 2 & 1 \end{pmatrix} =$$

result of applying $\begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & 1 & 4 \end{pmatrix}$ after applying $\begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 3 & 2 & 1 \end{pmatrix}$ or $n \begin{pmatrix} 1 & 2 & 3 & 4 \\ 4 & 1 & 3 & 2 \end{pmatrix}$.

Let f_i, g_i be maps. It isn't hard to verify that the Koszul convention implies that $(f_1 \otimes g_1) \circ (f_2 \otimes g_2) = (-1)^{\deg(f_2) \cdot \deg(g_1)} (f_1 \circ f_2 \otimes g_1 \circ g_2)$.

We will also follow the convention that, if f is a map between chain-complexes, $\partial f = \partial \circ f - (-1)^{\deg(f)} f \circ \partial$. The compositions of a map with boundary operations will be denoted by $\partial \circ f$ and $f \circ \partial$ — see [1]. This convention clearly implies that $\partial(f \circ g) = (\partial f) \circ g + (-1)^{\deg(f)} f \circ (\partial g)$. We will call any map f with $\partial f = 0$ a chain-map. We will also follow the convention that if C is a chain-complex and $\uparrow: C \rightarrow \Sigma C$ and $\downarrow: C \rightarrow \Sigma^{-1}C$ are, respectively, the suspension and desuspension maps, then $\uparrow$ and $\downarrow$ are both chain-maps. This implies that the boundary of ΣC is $-\uparrow \circ \partial_C \circ \downarrow$ and the boundary of $\Sigma^{-1}C$ is $-\downarrow \circ \partial_C \circ \uparrow$.

Definition 2.5. We will use the symbol T to denote the transposition operator for tensor products of chain-complexes $T: C \otimes D \rightarrow D \otimes C$, where $T(c \otimes d) = (-1)^{\dim(c) \cdot \dim(d)} d \otimes c$.

Proposition 2.6. Let C and D be chain-complexes. Then there exist isomorphisms

$$L_k = \downarrow_{C \otimes D}^k \circ (\uparrow_C^k \otimes 1_D): \Sigma^{-k} C \otimes D \rightarrow \Sigma^{-k}(C \otimes D)$$

sending $c \otimes d \in \Sigma^{-k} C_i \otimes D_j$ to $c \otimes d \in \Sigma^{-k}(C \otimes D)_{i+j}$, $r, d \in D_j$, and

$$M_k = \downarrow_{C \otimes D}^k \circ (1_C \otimes \uparrow_D^k): C \otimes \Sigma^{-k} D \rightarrow \Sigma^{-k}(C \otimes D)$$

sending $c \otimes d \in C_i \otimes \Sigma^{-k} D_j$ to $(-1)^{ik} c \otimes d \in \Sigma^{-k}(C \otimes D)_{i+j}$, for $c \in C_i$ and $d \in \Sigma^{-k} D_j = D_{j+k}$.

Definition 2.7. Let $\alpha_i, i = 1, \dots, n$ be a sequence of nonnegative integers whose sum is $|\alpha|$. Define a set-mapping symmetric groups

$$T_{\alpha_1, \dots, \alpha_n}: S_n \rightarrow S_{|\alpha|}$$

as follows:

1. for i between 1 and n , let L_i denote the length- α_i integer sequence:
2. , where $A_i = \sum_{j=1}^{i-1} \alpha_j$ — so, for instance, the concatenation of all of the L_i is the sequence of integers from 1 to $|\alpha|$;
3. $T_{\alpha_1, \dots, \alpha_n}(\sigma)$ is the permutation on the integers $1, \dots, |\alpha|$ that permutes the blocks $\{L_i\}$ via σ . In other words, σ is the permutation

$$\begin{pmatrix} 1 & \dots & n \\ \sigma(1) & \dots & \sigma(n) \end{pmatrix}$$

4

then $T_{\alpha_1, \dots, \alpha_n}(\sigma)$ is the permutation defined by writing

$$\begin{pmatrix} L_1 & \dots & L_n \\ L_{\sigma(1)} & \dots & L_{\sigma(n)} \end{pmatrix}$$

and regarding the upper and lower rows as sequences length $|\alpha|$.

Remark 2.8. Do not confuse the T -maps defined here with the transposition map for tensor products of chain-complexes. We will use the special notation T_i to represent $T_{1, \dots, 2, \dots, 1}$, where the 2 occurs in the i^{th} position. The two notations don't conflict since the old notation is never used in the case when $n = 1$. Here is an example of the computation of $T_{2,1,3}((1, 3, 2)) = T_{2,1,3} \begin{pmatrix} 1 & 2 & 3 \\ 3 & 1 & 2 \end{pmatrix} : L_1 = \{1\}2, L_2 = \{3\}, L_3 = \{4, 5, 6\}$. The permutation maps the ordered set $\{1, 2, 3\}$ to $\{3, 1, 2\}$, so we carry out the corresponding mapping of the sequences $\{L_1, L_2, L_3\}$ to get $\begin{pmatrix} L_1 & L_2 & L_3 \\ L_3 & L_1 & L_2 \end{pmatrix} = \begin{pmatrix} \{1, 2\} & \{3\} & \{4, 5, 6\} \\ \{4, 5, 6\} & \{1, 2\} & \{3\} \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 \\ 4 & 5 & 6 & 1 & 2 & 3 \end{pmatrix}$ (or $((1, 4)(2, 5)(3, 6))$, in cycle notation).

Definition 2.9. A sequence of differential graded $\mathbb{Z}$ -free modules, $\{\mathcal{U}_i\}$, will be said to form an *operad* if they satisfy the following conditions:

1. there exists a *unit map* (defined by the commutative diagrams below)

$$\eta: \mathbb{Z} \rightarrow \mathcal{U}_1$$

2. for all $i > 1$, $\mathcal{U}_i$ is equipped with a left action of S_i , the symmetric group.
3. for all $k \geq 1$, and $i_s \geq 0$ there are maps

$$\gamma: \mathcal{U}_{i_1} \otimes \dots \otimes \mathcal{U}_{i_k} \otimes \mathcal{U}_k \rightarrow \mathcal{U}_i$$

where $i = \sum_{j=1}^k i_j$.

The γ -maps must satisfy the following conditions:

Associativity: the following diagrams commute, where $\sum j_t = j$, $\sum i_s = i$, and $g_\alpha = \sum_{\ell=1}^\alpha j_\ell$ and $h_s = \sum_{\beta=g_{s-1}+1}^{g_s} i_\beta$:

$$\begin{array}{ccc} \left(\bigotimes_{s=1}^j \mathcal{U}_{i_s} \right) \otimes \left(\bigotimes_{t=1}^k \mathcal{U}_{j_t} \right) \otimes \mathcal{U}_k & \xrightarrow{\text{Id} \otimes \gamma} & \left(\bigotimes_{s=1}^j \mathcal{U}_{i_s} \right) \otimes \mathcal{U}_j \\ \downarrow \text{shuffle} & & \downarrow \gamma \\ \left(\left(\bigotimes_{q=1}^{j_t} \mathcal{U}_{i_{g_{t-1}+q}} \right) \otimes \bigotimes_{t=1}^k \mathcal{U}_{j_t} \right) \otimes \mathcal{U}_k & \xrightarrow{(\otimes_t \gamma) \otimes \text{Id}} & \left(\bigotimes_{t=1}^k \mathcal{U}_{h_k} \right) \otimes \mathcal{U}_k \\ & & \uparrow \gamma \\ & & \mathcal{U}_i \end{array}$$

Units: the following diagrams commute:

$$\begin{array}{ccc} \mathbb{Z}^k \otimes \mathcal{U}_k & \xrightarrow{\cong} & \mathcal{U}_k \\ \eta^k \otimes \text{Id} \downarrow & \nearrow \gamma & \\ \mathcal{U}_1^k \otimes \mathcal{U}_k & & \end{array} \quad \begin{array}{ccc} \mathcal{U}_k \otimes \mathbb{Z} & \xrightarrow{\cong} & \mathcal{U}_k \\ \text{Id} \otimes \eta \downarrow & \nearrow \gamma & \\ \mathcal{U}_k \otimes \mathcal{U}_1 & & \end{array}$$

Equivariance: the following diagrams commute:

$$\begin{array}{ccc} \mathcal{U}_{j_1} \otimes \cdots \otimes \mathcal{U}_{j_k} \otimes \mathcal{U}_k & \xrightarrow{\gamma} & \mathcal{U}_j \\ \sigma^{-1} \otimes \sigma \downarrow & & \downarrow T_{j_1, \dots, j_k}(\sigma) \\ \mathcal{U}_{j_{\sigma(1)}} \otimes \cdots \otimes \mathcal{U}_{j_{\sigma(k)}} \otimes \mathcal{U}_k & \xrightarrow{\gamma} & \mathcal{U}_j \end{array}$$

where $\sigma \in S_k$, and the σ^{-1} on the left permutes the factors $\{\mathcal{U}_{j_i}\}$ and the σ on the right simply acts on $\mathcal{U}_k$. See 2.7 for a definition of $T_{j_1, \dots, j_k}(\sigma)$.

$$\begin{array}{ccc} \mathcal{U}_{j_1} \otimes \cdots \otimes \mathcal{U}_{j_k} \otimes \mathcal{U}_k & \xrightarrow{\gamma} & \mathcal{U}_j \\ \tau_1 \otimes \cdots \otimes \tau_k \otimes \text{Id} \downarrow & & \downarrow \tau_1 \oplus \cdots \oplus \tau_k \\ \mathcal{U}_{j_{\sigma(1)}} \otimes \cdots \otimes \mathcal{U}_{j_{\sigma(k)}} \otimes \mathcal{U}_k & \xrightarrow{\gamma} & \mathcal{U}_j \end{array}$$

where $\tau_s \in S_{j_s}$ and $\tau_1 \oplus \cdots \oplus \tau_k \in S_j$ is the block sum.

Remark 2.10. The alert reader will notice a discrepancy between our definition of operad and that in [2] (on which it was based). The difference is due to our using operads as parameters for systems of *maps*, rather than *n*-ary operations. We, consequently, compose elements of an operad as one composes *maps*, i.e. the second operand is to the *left* of the first. This is also why the symmetric groups act on the *left* rather than on the right.

We will frequently want to think of operads in other terms:

Definition 2.11. Let $\mathcal{U}$ be an operad as defined above. Given $k_1 \geq k_2 > 0$, define the i^{th} *composition*

$$\circ_i: \mathcal{U}_{k_2} \otimes \mathcal{U}_{k_1} \rightarrow \mathcal{U}_{k_1+k_2}$$

as the composite

$$(2.1) \quad \underbrace{\mathbb{Z} \otimes \cdots \otimes \mathbb{Z} \otimes \mathcal{U}_{k_2} \otimes \mathbb{Z} \otimes \cdots \otimes \mathbb{Z} \otimes \mathcal{U}_{k_1}}_{i^{\text{th}} \text{ factor}} \rightarrow \underbrace{\mathcal{U}_1 \otimes \cdots \otimes \mathcal{U}_1 \otimes \mathcal{U}_{k_2} \otimes \mathcal{U}_1 \otimes \cdots \otimes \mathcal{U}_1}_{i^{\text{th}} \text{ factor}} \otimes \mathcal{U}_{k_1} \rightarrow \mathcal{U}_{k_1+k_2-1}$$

where the final map on the right is γ .

These compositions satisfy the following conditions, for all $a \in \mathcal{U}_n$, $b \in \mathcal{U}_m$, and $c \in \mathcal{U}_t$:

Associativity: $(a \circ_i b) \circ_j c = a \circ_{i+j-1} (b \circ_j c)$

Commutativity: $a \circ_{i+m-1} (b \circ_j c) = (-1)^{mn} b \circ_j (a \circ_i c)$

Equivariance: $a \circ_{\sigma(i)} (\sigma \cdot b) = \underbrace{T_{1, \dots, n, \dots, 1}(\sigma)}_{i^{\text{th}} \text{ position}} \cdot (a \circ_i b)$

Remark 2.12. In [6], I originally *defined* operads (or formal coalgebras) in terms of these compositions. It turned out that I'd recapitulated the historical sequence of events: operads were originally defined this way and called *composition algebras*. I am indebted to Jim Stasheff for pointing this out to me.

Given this definition of operad, we recover the γ map in 2.9 by setting:

$$\gamma(u_{i_1} \otimes \cdots \otimes u_{i_k}) = u_{i_1} \circ_1 \cdots \circ_{k-1} u_{i_k} \circ_k u_k$$

(where the implied parentheses associate to the right). It is left to the reader to verify that the two definitions are equivalent (the commutativity condition, here, is a special case of the equivariance condition).

Morphisms of operads are defined in the obvious way:

Definition 2.13. Given two operads $\mathcal{U}$ and $\mathcal{V}$, a *morphism*

$$f: \mathcal{U} \rightarrow \mathcal{V}$$

is a sequence of chain-maps

$$f_i: \mathcal{U}_i \rightarrow \mathcal{V}_i$$

commuting with all the diagrams in 2.9 or (equivalently) preserving the composition operations in 2.11.

Now we give some examples:

Definition 2.14. The operad $\mathfrak{S}_0$ is defined via

1. Its n^{th} component is $\mathbb{Z}S_n$ — a chain-complex concentrated in dimension 0.
2. The composition operations are defined by

$$\gamma(\sigma_{i_1} \otimes \cdots \otimes \sigma_{i_k} \otimes \sigma) = \sigma_{i_{\sigma(k)}} \oplus \cdots \oplus \sigma_{i_{\sigma(k)}} \circ T_{i_1, \dots, i_k}(\sigma)$$

Remark 2.15. This was denoted $\mathcal{M}$ in [2].

Verification that this satisfies the required identities is left to the reader as an exercise.

Definition 2.16. Let $\mathfrak{S}$ denote the E_∞ -operad with components

$$\mathbb{R}S_n$$

— the bar resolutions of $\mathbb{Z}$ over $\mathbb{Z}S_n$ for all $n > 0$. This is an important operad and [5, § 2.3] is devoted to it.

Remark 2.17. This is the result of applying the “unreduced bar construction” to the previous example.

Definition 2.18. Coassoc is an operad defined to have one basis element $\{b_i\}$ for all integers $i \geq 0$. Here the rank of b_i is i and the degree is 0 and these elements satisfy the composition-law: $b_i \circ_\alpha b_j = b_{i+j-1}$ regardless of the value of α , which can run from 1 to j . The differential of this operad is identically zero.

Now we define two important operads associated to any $\mathbb{Z}$ -module.

Definition 2.19. Let C be a DGA-module with augmentation $\epsilon: C \rightarrow \mathbb{Z}$, and with the property that $C_0 = \mathbb{Z}$. Then the *Coendomorphism* operad, $\text{CoEnd}(C)$, is defined to be the operad with:

1. component of rank $i = \text{Hom}_{\mathbb{Z}}(C, C^i)$, with the differential induced by that of C and C^i . The dimension of an element of $\text{Hom}_{\mathbb{Z}}(C, C^i)$ (for some i) is defined to be its degree as a map.
2. The $\mathbb{Z}$ -summand is generated by one element, e , of rank 0.

Remark 2.20. Both operads are unitary — their common identity element is the identity map $\text{id} \in \text{Hom}_{\mathbb{Z}}(C, C)$. One motivation for operads is that they model the iterated coproducts that occur in $\text{CoEnd}(\ast)$. We will use operads as an algebraic framework for defining other constructs that have topological applications.

In like fashion, we define the *endomorphism operad*:

Definition 2.21. If C is a DGA-module, the *endomorphism operad*, $\text{End}(C)$ is defined to have components

$$\text{Hom}_{\mathbb{Z}}(C^n, C)$$

and compositions that coincide with endomorphism compositions.

Now we consider a special class of operads that play a crucial role in the sequel:

Definition 2.22. An operad $\mathcal{U} = \{\mathcal{U}_n\}$ will be called an E_{∞} -operad if $\mathcal{U}_n$ is a $\mathbb{Z}S_n$ -free resolution of $\mathbb{Z}$ for all $n > 0$.

Given these definitions, we can define:

Definition 2.23. Let $\mathcal{U}$ be an operad and let C be a DG-module equipped with a morphism (of operads)

$$f: \mathcal{U} \rightarrow \text{CoEnd}(C)$$

Then C is called a *coalgebra* over $\mathcal{U}$ with structure map f . If C is equipped with a morphism of operads

$$f: \mathcal{U} \rightarrow \text{End}(C)$$

then C is called a *algebra* over $\mathcal{U}$ with structure map f .

Remark 2.24. A coalgebra, C , over an operad, $\mathcal{U}$, is a sequence of maps

$$f_n: \mathcal{U} \otimes C \rightarrow C^n$$

for all $n > 0$, where f_n is $\mathbb{Z}S_n$ -equivariant. These maps are related in the sense that they fit into commutative diagrams:

$$\begin{array}{ccc} \mathcal{U}_n \otimes \mathcal{U}_m \otimes C & \xlongequal{\quad} & \mathcal{U}_n \otimes \mathcal{U}_m \otimes C \\ \downarrow 1 \otimes f_m & & \downarrow \circ_i \\ \mathcal{U}_n \otimes C^m & & \mathcal{U}_{n+m-1} \otimes C \\ \downarrow V_{i-1} & & \downarrow f_{n+m-1} \\ C^{i-1} \otimes \mathcal{U}_n \otimes C \otimes C^{m-i} & \xrightarrow{1 \otimes \cdots \otimes f_n \otimes \cdots \otimes 1} & C^{m+1} \end{array}$$

for all $n, m \geq 1$ and $1 \leq i \leq m$. Here $V: \mathcal{U}_n \otimes C^m \rightarrow C^{i-1} \otimes \mathcal{U}_n \otimes C \otimes C^{m-i}$ is the map that shuffles the factor $\mathcal{U}_n$ to the right of $i-1$ factors of C . In other words: The abstract composition-operations in $\mathcal{U}$ exactly correspond to compositions of maps in $\{\text{Hom}_{\mathbb{Z}}(C, C^n)\}$. We exploit this behavior in applications of coalgebras over operads, using an explicit knowledge of the algebraic structure of $\mathcal{U}$.

In very simple cases, one can explicitly describe the maps defining a coalgebra over an operad:

Definition 2.25. Define the coalgebra, I — *the unit interval* — over $\mathfrak{S}$ via:

1. Its $\mathbb{Z}$ -generators are $\{p_0, p_1\}$ in dimension 0 and q in dimension 1, and its adjoint structure map is $r_n: RS_n \otimes I \rightarrow I^n$, for all $n > 1$.
2. The coproduct is given by $r_2([] \otimes p_i) = p_i \otimes p_i$, $i = 0, 1$, and $r_2([] \otimes q) = p_0 \otimes q + q \otimes p_1$.
3. The higher coproducts are given by $r_2([(1, 2)] \otimes q) = q \otimes q$, $r_2([(1, 2)] \otimes p_i) = 0$, $i = 0, 1$ and $r_2(a \otimes I) = 0$, where $a \in RS_2$ has dimension > 1 .

Remark 2.26. This, coupled with the operad-identities in $\mathfrak{S}$ suffice to define the coalgebra structure of I in all dimensions and for all degrees.

Proposition 2.27. *Coassociative coalgebras are precisely the coalgebras over Coassoc.*

Remark 2.28. There are some subtleties to this definition, however. It is valid if we regard Coassoc as a non- Σ operad. If we regard it as an operad with *trivial* symmetric group action, then we have defined coassociative, *cocommutative* coalgebras.

We can define tensor products of operads:

Definition 2.29. Let $\mathcal{U}_1$ and $\mathcal{U}_2$ be operads. Then $\mathcal{U}_1 \otimes \mathcal{U}_2$ is defined to have:

1. component of rank $i = (\mathcal{U}_1)_i \otimes (\mathcal{U}_2)_i$, where $(\mathcal{U}_1)_i$ and $(\mathcal{U}_2)_i$ are, respectively, the components of rank i of $\mathcal{U}_1$ and $\mathcal{U}_2$;
2. composition operations defined via $(a \otimes b) \circ_i (c \otimes d) = (-1)^{\dim(b) \dim(c)} (a \circ_i c) \otimes (b \circ_i d)$, for $a, c \in \mathcal{U}_1$ and $b, d \in \mathcal{U}_2$.

We conclude this section with

Definition 2.30. An operad, $\mathfrak{R}$, will be called an *operad-coalgebra* if there exists a co-associative morphism of operads

$$\Delta: \mathfrak{R} \rightarrow \mathfrak{R} \otimes \mathfrak{R}$$

Remark 2.31. Operad-coalgebras are important in certain homotopy-theoretic contexts — for instance in the study of the bar and cobar constructions. The $\mathfrak{S}$ -operad is a particularly important operad of this type.

3. SUSPENSIONS

3.1. Chain-complexes and operads. In this section, we will compute the effect of a suspension in some detail. We begin with a word on notation. There is the well-known concept of suspension of a chain-complex: this is just the given chain-complex with dimensions shifted up or down. In [5], we defined the concept of suspension of an m -coalgebra that corresponded to the topological notion of suspension. Now the problem: although m -coalgebras are also chain-complexes, their suspensions as *chain-complexes* are *not* the same as their suspensions as *m -coalgebras*.

We will consequently use the notation SC for the suspension of C as an *m -coalgebra* (this is the suspension that has topological significance).

Definition 3.1. Let $C = \Sigma\mathbb{Z}$ be the chain-complex concentrated in dimension 1 and define $\text{Susp} = \text{CoEnd}(C)$, the *suspension operad*. In like fashion, we can define $\text{Susp}^{-1} = \text{CoEnd}(\Sigma^{-1}\mathbb{Z})$, the *desuspension operad*.

Remark 3.2. Note that Susp_n is a chain-complex concentrated in dimension $n-1$ equal to $\Sigma^{n-1}\mathbb{Z}$, with compositions defined by

$$s_n \circ_i s_m = (-1)^{(i-1)(n-1)} s_{n+m-1}$$

where $s_n \in \text{Susp}_n$ is the canonical generator. The S_n -action on s_n is defined by

$$\sigma \cdot s_n = (-1)^{\text{parity}(\sigma)} s_n$$

where $\sigma \in S_n$.

Note that there are two distinct, but canonically isomorphic, ways of defining Susp . We have selected the definition that corresponds to composing maps $f, g \in \text{CoEnd}(C)$ to have $f \circ_i g$ mean “the map g followed by the map f .”

This definition was motivated by

Proposition 3.3. *If C is a chain-complex*

$$\text{CoEnd}(\Sigma C) = \text{Susp} \otimes \text{CoEnd}(C)$$

Remark 3.4. Note that the suspensions here are those of *chain-complexes* (i.e., simple dimension-shifting) rather than the (topologically motivated) notion of *m-coalgebra* suspension defined in [5, 3.36]. We will connect these two notions later in this section.

For the component of degree n , the isomorphisms in question are precisely $\text{Hom}_{\mathbb{Z}}(\downarrow, \underbrace{\downarrow \otimes \cdots \otimes \downarrow}_{n \text{ times}})$, which lower dimension by $n-1$.

Proof. Note that $\Sigma C = \Sigma\mathbb{Z} \otimes C$. The conclusion follows from the injective operad-morphism (see [5, Proposition 3.4])

$$\mathfrak{E}: \text{CoEnd}(\Sigma\mathbb{Z}) \otimes \text{CoEnd}(C) \rightarrow \text{CoEnd}(\Sigma\mathbb{Z} \otimes C)$$

which is an isomorphism in this case (since $D = \text{Susp}$ has finitely generated components). $\square$

We can use this to define the concept of *suspension of an operad*:

Definition 3.5. Given an operad A , we define its suspension ΣA to be the operad $\text{Susp} \otimes A$ and its *desuspension*, $\Sigma^{-1}A$, to be the operad $\text{Susp}^{-1} \otimes A$. The *suspension isomorphism* $\mathcal{J}: A \rightarrow \Sigma A$ is a set of isomorphisms

$$\mathcal{J}_n: A_n \rightarrow (\Sigma A)_n$$

of degree $n-1$ (i.e., $\mathcal{J}_n$ raises dimension by $n-1$) for all $n > 0$ such that:

1. The identity

$$\mathcal{J}_n(g \cdot x) = (-1)^{\text{parity}(g)} g \cdot \mathcal{J}_n(x)$$

holds for all $n > 0$, all $x \in A_n$ and all $g \in S_n$.

2. The identity

$$\mathcal{J}_{n+m-1}(a \circ_i b) = (-1)^{(m-1)\dim a + (n-1)(i-1)} \mathcal{J}_n(a) \circ_i \mathcal{J}_m(b)$$

holds for all $n, m > 0$, all $a \in A_n$ and $b \in A_m$, and all $0 \leq i \leq m$.

Although suspension-isomorphisms are not operad-morphisms (they do not even preserve dimension) they clearly define a functor from the category of operads to itself.

Definition 3.6. Let $k \geq 0$ be an integer and define $\Sigma^{-k}\mathfrak{L}$ to be the category of DG-coalgebras over the operad $\Sigma^{-k}\mathfrak{S}$ that are k -fold desuspensions (in the Σ -sense) of $\mathfrak{m}$ -coalgebras. Define $\Sigma^{-k}\mathfrak{L} = \Sigma^{-k}\mathfrak{L}_0[H^{-1}]$ where H is the class of morphisms of coalgebras over $\Sigma^{-k}\mathfrak{S}$ inducing *homology isomorphisms* of underlying chain-complexes.

By *definition*

Proposition 3.7. *Desuspension defines isomorphisms of categories*

$$\downarrow^k: \mathfrak{L} \rightarrow \Sigma^{-k}\mathfrak{L}$$

for all $k \geq 0$.

3.2. $\mathfrak{m}$ -coalgebras. For the remainder of this section, we will focus on $\mathfrak{m}$ -coalgebras over the operad $\mathfrak{S}$ and consider suspensions of $\mathfrak{m}$ -coalgebras as defined in [5, 3.36].

There are important differences between suspending $\mathfrak{m}$ -coalgebras and arbitrary coalgebras over an operad. For instance $\mathfrak{m}$ -coalgebras contain a sub-coalgebra isomorphic to the *trivial $\mathfrak{m}$ -coalgebra*, $\mathbb{Z}$ — the *basepoint*. In addition, there exists a canonical surjection of coalgebras

$$C \rightarrow \mathbb{Z}$$

that is a left-inverse to inclusion of the basepoint. The basepoint does *not* participate in suspension.

Definition 3.8. Let k be an integer ≥ 0 , and let C be an object of $\Sigma^{-k}\mathfrak{L}_0$. Then we define $C^+ = C/\Sigma^{-k}\mathbb{Z}$, where $\Sigma^{-k}\mathbb{Z} = C_{-k}$ is the (desuspended) basepoint.

Remark 3.9. By [5, 2.35], this quotient is a coalgebra over $\Sigma^{-k}\mathfrak{S}$.

Definition 3.10. Let k be an integer ≥ 0 , and let C be an object of $\Sigma^{-k}\mathfrak{L}_0$ with structure-map

$$a: \Sigma^{-k}\mathfrak{S} \rightarrow \text{CoEnd}(C)$$

We will call C *reduced* if its coproduct is trivial — i.e.

$$\Delta(x) = a(\Sigma^{-k}[])(x) = \Sigma^{-k}1 \otimes x + x \otimes \Sigma^{-k}1$$

for all $x \in C$, where $1 \in C_{-k}$ is the (desuspended) basepoint.

The following is immediate:

Proposition 3.11. *Let k be an integer ≥ 0 , and let C and D be reduced objects of $\Sigma^{-k}\mathfrak{L}_0$. Then $C \cong D$ if and only if $C^+ \cong D^+$.*

Now we are in a position to begin to calculate the $S \wedge *$ -suspension of $\mathfrak{m}$ -coalgebras.

Proposition 3.12. *Let $[0, 1]$ denote the standard 1-simplex, with vertices $\{[0], [1]\}$, and let Susp be the operad defined in 3.1. Then there exists a surjective operad-morphism*

$$(3.1) \quad \tau: \text{CoEnd}(\mathcal{C}([0, 1])) \rightarrow \text{Susp}$$

Proof. This is just the morphism of $\mathrm{CoEnd}(\ast)$ -operads induced by the chain-map

$$\mathcal{C}([0, 1]) \rightarrow \mathcal{C}([0, 1])/\mathcal{C}([0, 1])_0 = \Sigma\mathbb{Z}$$

□

Proposition 3.13. *There exists a surjective operad-morphism*

$$\mathfrak{V}: \mathfrak{S} \rightarrow \Sigma\mathfrak{S}$$

that makes the following diagram commute

$$\begin{array}{ccc} & \Sigma\mathfrak{S} & \\ \mathfrak{V} \nearrow & & \searrow \mathfrak{J} \otimes 1 \circ \Delta \circ \mathfrak{J}^{-1} \\ \mathfrak{S} & & \Sigma\mathfrak{S} \otimes \mathfrak{S} \\ \Delta \searrow & & \nearrow \mathfrak{V} \otimes 1 \\ & \mathfrak{S} \otimes \mathfrak{S} & \end{array}$$

Remark 3.14. The morphism $\mathfrak{V}$ coincides with the *determinant map* defined in [5]. Note that [5, Proposition 2.26] does not apply since $\Sigma\mathfrak{S}$ and $\Sigma^{-1}\mathfrak{S}$ are not E_∞ -operads. Although the $\{\mathfrak{V}_n\}$ do not induce homology isomorphisms, they do define cohomology classes

$$\{\alpha_n \in H^{n-1}(S_n, \mathbb{Z})\}$$

for all $n > 0$.

Proof. Identify $\Sigma\mathfrak{S}$ with the operad $\mathrm{Susp} \otimes \mathfrak{S}$, by 3.5. Now form the composite

$$(3.2) \quad \mathfrak{S} \xrightarrow{\Delta} \mathfrak{S} \otimes \mathfrak{S} \xrightarrow{u \otimes 1} \mathrm{CoEnd}(U) \otimes \mathfrak{S} \xrightarrow{\tau \otimes 1} \mathrm{Susp} \otimes \mathfrak{S} = \Sigma\mathfrak{S}$$

where

1. $\Delta: \mathfrak{S} \rightarrow \mathfrak{S} \otimes \mathfrak{S}$ is the canonical coproduct defined in [5],
2. the morphism τ is defined in 3.1,
3. $U = \mathcal{C}([0, 1])$,
4. $u: \mathfrak{S} \rightarrow \mathrm{CoEnd}(U)$ is the structure map of the $\mathfrak{m}$ -structure of $\mathcal{C}([0, 1])$.

□

We will use this to describe the $\mathfrak{m}$ -structure of suspensions.

Proposition 3.15. *Let C be an $\mathfrak{m}$ -coalgebra over $\mathfrak{S}$. Then*

$$\Sigma C^+ = (SC)^+$$

as chain-complexes.

Proof. Recall that

$$SC = \mathcal{C}(S^1) \wedge C = \mathcal{C}(I) \otimes C / ([0] \otimes C^+ + [1] \otimes C^+ \otimes + [0, 1]^+ \otimes p)$$

where $p \in C_0 = \mathbb{Z}$ is the basepoint and $[0]$ is the basepoint of $[0, 1]$. This means that the basepoint of SC is $[0] \otimes p$ and $(SC)^+ = [0, 1] \otimes C^+ = \Sigma\mathbb{Z} \otimes C^+ = \Sigma C^+$. □

The following result relates our two distinct notions of suspension

JUSTIN R. SMITH

Theorem 3.16. *Let C be an m -coalgebra over $\mathfrak{S}$ with structure map*

$$a_C: \mathfrak{S} \rightarrow \text{CoEnd}(C)$$

whose suspension (as an m -coalgebra) is ΣC with structure map

$$a_{\Sigma C}: \mathfrak{S} \rightarrow \text{CoEnd}(\Sigma C)$$

Then $\Sigma C^+ = (SC)^+$. Suppose that the corresponding reduced m -coalgebras have the structure maps

$$a_C^+: \mathfrak{S} \rightarrow \text{CoEnd}(C^+)$$

and

$$a_{\Sigma C}^+: \mathfrak{S} \rightarrow \text{CoEnd}((SC)^+) = \text{CoEnd}(\Sigma C^+)$$

respectively. Then the following diagram commutes

$$(3.3) \quad \begin{array}{ccccc} \text{CoEnd}((SC)^+) & \xlongequal{\quad} & \text{CoEnd}(\Sigma C^+) & \xleftarrow{\mathfrak{J}} & \text{CoEnd}(C^+) \\ \uparrow a_{\Sigma C}^+ & & \uparrow \Sigma a_C^+ & & \uparrow a_C^+ \\ \mathfrak{S} & \xrightarrow{\mathfrak{S}} & \Sigma \mathfrak{S} & \xleftarrow{\mathfrak{J}} & \mathfrak{S} \end{array}$$

where all vertical maps are operad-morphisms and the $\mathfrak{J}$ -maps are suspension-isomorphisms of operads (see 3.5).

Proof. This follows from 3.13 and the definition of the m -structure of the (geometric) suspension. Let C be an m -coalgebra over $\mathfrak{S}$ with structure map

$$a: \mathfrak{S} \rightarrow \text{CoEnd}(C)$$

and let the structure map of the unit interval $I = \mathcal{C}([0, 1])$ be

$$u: \mathfrak{S} \rightarrow \text{CoEnd}(I)$$

Then the suspension has a structure map that is the composite

$$(3.4) \quad \begin{array}{ccc} \mathfrak{S} & \xlongequal{\quad} & \mathfrak{S} \\ \Delta \downarrow & & \downarrow \Delta \\ \mathfrak{S} \otimes \mathfrak{S} & \xlongequal{\quad} & \mathfrak{S} \otimes \mathfrak{S} \\ u \otimes 1 \downarrow & & \downarrow u \otimes a \\ \text{CoEnd}(I) \otimes \mathfrak{S} & \xrightarrow{1 \otimes a} & \text{CoEnd}(I) \otimes \text{CoEnd}(C) \\ \parallel & & \downarrow \mathfrak{e} \\ \text{CoEnd}(I) \otimes \mathfrak{S} & \xrightarrow{\mathfrak{e} \circ 1 \otimes a} & \text{CoEnd}(I \otimes C) \\ \{\tau \otimes 1\} \downarrow & & \downarrow q \\ \text{Susp} \otimes \mathfrak{S} & \xrightarrow{(\mathfrak{e} \circ 1 \otimes a)^*} & \text{CoEnd}((SC)^+) \\ \parallel & & \parallel \\ \Sigma \mathfrak{S} & \xrightarrow{\Sigma a^+} & \text{CoEnd}(\Sigma C^+) \end{array}$$

Where

1. $\mathfrak{E}: \text{CoEnd}(I) \otimes \text{CoEnd}(C) \rightarrow \text{CoEnd}(I \otimes C)$ is the operad morphism defined in [5, Proposition 3.4]
2. $q: \text{CoEnd}(I \otimes C) \rightarrow \text{CoEnd}((SC)^+)$ is induced by the quotient map $I \otimes C \rightarrow I \otimes C / ([0] \otimes C^+ + [1] \otimes C^+ + [0, 1]^+ \otimes p + [0] \otimes p) = (SC)^+$

Now observe that the left column is $\mathfrak{V}$ and that the right is the structure map of $(SC)^+$.

□

Corollary 3.17. *Let C and D be m -coalgebras over $\mathfrak{S}$ with reduced structure morphisms*

$$a_C: \mathfrak{S} \rightarrow \text{CoEnd}(C^+)$$

and

$$a_D: \mathfrak{S} \rightarrow \text{CoEnd}(D^+)$$

respectively. Then a morphism $f: SC \rightarrow D$ gives rise to a commutative diagram

$$(3.5) \quad \begin{array}{ccc} C^+ & \xrightarrow{\Sigma^{-1}f^+} & \Sigma^{-1}D^+ \\ a_C \uparrow & & \uparrow \Sigma^{-1}a_D \\ \mathfrak{S} & \xleftarrow{\mathfrak{u}} & \Sigma^{-1}\mathfrak{S} \end{array}$$

Proof. Just desuspend the left square of diagram 3.3 (and interchange its columns) to get

$$\begin{array}{ccc} \text{CoEnd}(C^+) & \xlongequal{\quad} & \Sigma^{-1}\text{CoEnd}((SC)^+) \\ a_C^+ \uparrow & & \uparrow \Sigma^{-1}a_{SC}^+ \\ \mathfrak{S} & \xleftarrow{\mathfrak{u}} & \Sigma^{-1}\mathfrak{S} \end{array}$$

and splice in the commutative square representing a morphism $\Sigma^{-1}f: \Sigma^{-1}SC \rightarrow \Sigma^{-1}D$ (in $\mathfrak{L}_0^{-1}$)

$$\begin{array}{ccc} \Sigma^{-1}\text{CoEnd}((SC)^+) & \xrightarrow{\Sigma^{-1}f} & \Sigma^{-1}D^+ \\ \Sigma^{-1}a_{SC}^+ \uparrow & & \uparrow \Sigma^{-1}a_D \\ \Sigma^{-1}\mathfrak{S} & \xlongequal{\quad} & \Sigma^{-1}\mathfrak{S} \end{array}$$

The “outer rim” of the result is precisely 3.5.

□

The following is straightforward, but should be said

Definition 3.18. Given a coalgebra C over an operad $\mathfrak{R}_1$ with structure map

$$a: \mathfrak{R}_1 \rightarrow \text{CoEnd}(C)$$

and a morphism of operads

$$f: \mathfrak{R}_2 \rightarrow \mathfrak{R}_1$$

the *pullback*, f^*C is defined to be the coalgebra over $\mathfrak{R}_2$ with structure map

$$a \circ f: \mathfrak{R}_2 \rightarrow \text{CoEnd}(C)$$

We have two distinct ways of converting objects of $\mathfrak{L}$ into objects of $\Sigma^{-k}\mathfrak{L}$: by desuspending, as in 3.7, or by pulling back structure morphisms over

$$\mathfrak{U}^k = \Sigma^{-k}\mathfrak{V}^k: \Sigma^{-k}\mathfrak{S} \rightarrow \mathfrak{S}$$

as in 3.17.

The first method desuspends the underlying chain-complex (so we get a chain-complex that extends into negative dimensions), and the second does not. These two functors are related by:

Theorem 3.19. *Let C and D be m -coalgebras over $\mathfrak{S}$ with structure-maps a_C and a_D , respectively, let k be an integer > 0 . Then $S^k C$ is equivalent to $S^k D$ in $\mathfrak{L}$ if and only if $(\mathfrak{U}^k)^* C^+$ is equivalent to $(\mathfrak{U}^k)^* D^+$ in $\Sigma^{-k}\mathfrak{L}$.*

Proof. This follows by an inductive application of 3.16. □

Definition 3.20. Consider the inverse system of operads

$$\dots \rightarrow \Sigma^{-i}\mathfrak{S} \rightarrow \Sigma^{-i+1}\mathfrak{S} \rightarrow \dots \rightarrow \mathfrak{S}$$

whose maps are

$$\Sigma^{-i}\mathfrak{U}: \Sigma^{-i-1}\mathfrak{S} \rightarrow \Sigma^{-i}\mathfrak{S}$$

Denote the corresponding inverse limit by $\mathfrak{S}_{-\infty}$. The morphism $\mathfrak{U}: \Sigma^{-1}\mathfrak{S} \rightarrow \mathfrak{S}$ induces an operad-endomorphism

$$\mathfrak{U}: \mathfrak{S}_{-\infty} \rightarrow \mathfrak{S}_{-\infty}$$

In addition, $\mathfrak{S}_{-\infty}$ comes equipped with canonical operad-morphisms

$$\mathfrak{p}_n: \mathfrak{S}_{-\infty} \rightarrow \Sigma^{-n}\mathfrak{S}$$

Remark 3.21. We may think of $\mathfrak{S}_{-\infty}$ as “ $\Sigma^{-\infty}\mathfrak{S}$ ” and the projections $\mathfrak{p}_n$ as “ $\mathfrak{U}^{\infty-n}$ ”. This makes some sense if one notes that all of the $\mathfrak{U}^n$ maps are *surjective* so that there are isomorphisms

$$\Sigma^{-n}\mathfrak{S} \rightarrow F^n\{\Sigma^{-i}\mathfrak{S}\}$$

where $F^n\{\Sigma^{-i}\mathfrak{S}\}$ is a “finite” portion of the inverse limit — the sub-operad of

$$\prod_{i=0}^n \Sigma^{-i}\mathfrak{S} \subset \prod_{i=0}^{\infty} \Sigma^{-i}\mathfrak{S}$$

composed of sequences $\{\alpha_0, \dots, \alpha_n\}$ with $\alpha_i = \mathfrak{U}(\alpha_{i+1})$ for all $i < n$.

Definition 3.22. Let $\mathfrak{L}'_{-\infty}$ denote the category of coalgebras over $\mathfrak{S}_{-\infty}$ and let $\mathfrak{L}_{-\infty} = \mathfrak{L}'_{-\infty}[H^{-1}]$, where H is the class of morphisms inducing homology-isomorphisms.

Let $\Sigma^{-\infty}\mathfrak{L}_0 \subset \mathfrak{L}'_{-\infty}$ be the subcategory with the property that each object, $\mathcal{O}$, of $\Sigma^{-\infty}\mathfrak{L}_0$ is of the form $\mathfrak{p}_k^* Z$,

1. where $Z \in \Sigma^{-k}\mathfrak{L}_0$ for some value of k , and
2. $\mathfrak{p}_k: \mathfrak{S}_{-\infty} \rightarrow \Sigma^{-k}\mathfrak{S}$, as defined in 3.20.

Given an object $\mathcal{O}$, of $\Sigma^{-\infty}\mathfrak{L}_0$ of the form $\mathfrak{p}_k^*Z$, we will call the integer k the *level* of $\mathcal{O}$. In addition, define $\Sigma^{-\infty}\mathfrak{L} = \Sigma^{-\infty}\mathfrak{L}_0[H^{-1}]$, where H is the class of morphisms inducing homology isomorphisms. Then morphisms $m: \mathcal{O}_1 \rightarrow \mathcal{O}_2$ are diagrams of the form

$$\mathcal{O}_1 \leftrightarrow Z_1 \leftrightarrow \cdots \leftrightarrow Z_n \leftrightarrow \mathcal{O}_2$$

with each $\leftrightarrow$ representing a morphism to the left or right, and all left-pointing arrows inducing homology equivalences. We will define the *level of the morphism* m to be

$$\max\{\text{level}(\mathcal{O}_1), \text{level}(Z_1), \dots, \text{level}(Z_n), \text{level}(\mathcal{O}_2)\}$$

Corollary 3.23. *Let C and D be m -coalgebras. Then there exists an integer $k > 0$ and a morphism $S^k C \rightarrow S^k D$ in $\mathfrak{L}$ if and only if there exists a morphism $\mathfrak{p}_0^* C^+ \rightarrow \mathfrak{p}_0^* D^+$ in $\Sigma^{-\infty}\mathfrak{L}$. The two morphisms induce identical maps of chain-complexes. In particular, there exists an integer $k > 0$ such that $S^k C$ is equivalent to $S^k D$ in $\mathfrak{L}$ if and only if $\mathfrak{p}_0^* C^+$ is equivalent to $\mathfrak{p}_0^* D^+$ in $\Sigma^{-\infty}\mathfrak{L}$.*

Proof. We begin by proving the *if*-part: If there exists a morphism $S^k C \rightarrow S^k D$ then, by 3.19, $(\mathfrak{U}^k)^* C^+ \cong (S^k C)^+$ and $(\mathfrak{U}^k)^* D^+ \cong (S^k D)^+$ in $\Sigma^{-k}\mathfrak{L}$. Since $\mathfrak{p}_k \circ \mathfrak{U}^k = \mathfrak{p}_0$, the morphism $S^k C \rightarrow S^k D$ induces a morphism

$$\mathfrak{p}_0^* C^+ \rightarrow \mathfrak{p}_0^* D^+$$

in $\mathfrak{L}_{-\infty}$.

One the other hand, suppose there exists a morphism $\mathfrak{p}_0^* C^+ \rightarrow \mathfrak{p}_0^* D^+$ in $\Sigma^{-\infty}\mathfrak{L}$, of level k (see 3.22). Then there exists a sequence

$$(3.6) \quad \mathfrak{p}_0^* C^+ \leftrightarrow Z_1 \leftrightarrow \cdots \leftrightarrow Z_t \leftrightarrow \mathfrak{p}_0^* D^+$$

where each arrow, $\leftrightarrow$, points to the left or right and the left-pointing arrows induce homology isomorphisms. Suppose $Z_i = \mathfrak{p}_{k_i}^* W_i$, where $W_i \in \Sigma^{-k_i}\mathfrak{L}_0$. Then $k = \max\{k_i\}$. We convert 3.6 into a sequence

$$(\mathfrak{U}^k)^* C^+ \leftrightarrow (\mathfrak{U}^{k-k_1})^* Z_1 \leftrightarrow \cdots \leftrightarrow (\mathfrak{U}^{k-k_t})^* Z_t \leftrightarrow (\mathfrak{U}^k)^* D^+$$

which defines a morphism $(\mathfrak{U}^k)^* C^+ \rightarrow (\mathfrak{U}^k)^* D^+$ in $\Sigma^{-k}\mathfrak{L}$. The conclusion follows by 3.19. $\square$

Corollary 3.24. *Let X and Y be pointed, simply-connected semisimplicial sets. Then there exists an integer $k \geq 0$ and a pointed map of topological realizations $f: |S^k X| \rightarrow |S^k Y|$ if and only if there exists a morphism $g: \mathfrak{p}_0^* \mathcal{C}(X)^+ \rightarrow \mathfrak{p}_0^* \mathcal{C}(Y)^+$ in $\Sigma^{-\infty}\mathfrak{L}$. In this case, the following diagram commutes in $\mathfrak{L}$*

$$(3.7) \quad \begin{array}{ccc} \Sigma^k(\mathfrak{U}^k)^* \mathcal{C}(X)^+ & \xrightarrow{\Sigma^k g} & \Sigma^k(\mathfrak{U}^k)^* \mathcal{C}(Y)^+ \\ \parallel & & \parallel \\ \mathcal{C}(S^k X)^+ & & \mathcal{C}(S^k Y)^+ \\ \downarrow \iota_{S^k X} & & \downarrow \iota_{S^k Y} \\ \mathcal{C}(\dot{\Delta}(|S^k X|)) & \xrightarrow{\mathcal{C}(\dot{\Delta}(f))} & \mathcal{C}(\dot{\Delta}(|S^k Y|)) \end{array}$$

where

OPERADS AND ALGEBRAIC HOMOTOPY II:
SUSPENSIONS

JUSTIN R. SMITH

-
1. Σ^k denotes suspension of chain-complexes and base-operads. This is suspension in the sense of dimension-shifting.
 2. $\dot{\Delta}(\ast)$ denotes the 2-reduced singular complex functor.
 3. $|\ast|$ denotes topological realization functor.
 4. $\iota_X: \mathcal{C}(X) \rightarrow \mathcal{C}(\dot{\Delta}(|X|))$ is the canonical (injective) equivalence.

In particular X and Y are stably homotopy equivalent if and only if $\mathfrak{p}_0^* \mathcal{C}(X)^+$ and $\mathfrak{p}_0^* \mathcal{C}(Y)^+$ are equivalent in $\Sigma^{-\infty} \mathfrak{L}$.

Remark 3.25. Note that $\mathfrak{L}$ is the localized category $\mathfrak{L}_0[H^{-1}]$, so that the commutativity of 3.7 in $\mathfrak{L}$ implies its *homotopy* commutativity in $\mathfrak{L}_0$.

Note that the basepoints have fallen out of the picture: the stable homotopy type of X is determined by the reduced m-coalgebra $\mathcal{C}(X)^+$.

Proof. This follows from 3.23, which implies that $S^k \mathcal{C}(X)$ is equivalent to $S^k \mathcal{C}(Y)$ in $\mathfrak{L}$ and [5, Theorem 9.9]. □

REFERENCES

- [1] V. K. A. M. Gugenheim, *On a theorem of E. H. Brown*, Illinois J. of Math. **4** (1960), 292–311.
 - [2] I. Kriz and J. P. May, *Operads, algebras, modules and motives*, Astérisque, vol. 233, Société Mathématique de France, 1995.
 - [3] D. Quillen, *Homotopical algebra*, Lecture Notes in Mathematics, vol. 43, Springer-Verlag, 1967.
 - [4] ———, *Rational homotopy theory*, Ann. of Math. (2) **90** (1969), 205–295.
 - [5] Justin R. Smith, *Operads and Algebraic Homotopy*, preprint, March 2000. Can be downloaded from <http://vorpal.mcs.drexel.edu/research/current.shtml>.
 - [6] ———, *Iterating the cobar construction*, vol. 109, Memoirs of the A. M. S., no. 524, American Mathematical Society, Providence, Rhode Island, May 1994.
- E-mail address:* jsmith@mcs.drexel.edu
URL: <http://vorpal.mcs.drexel.edu>